

## C-CONFORMAL METRIC TRANSFORMATIONS ON FINSLERIAN HYPERSURFACE

S.K. NARASIMHAMURTHY<sup>1</sup>, PRADEEP KUMAR<sup>2</sup> AND C.S. BAGEWADI<sup>3</sup>

<sup>1</sup>Department of Mathematics, Kuvempu University,  
Shankaraghatta-577451, Shimoga, Karnataka, India, nmurthysk@gmail.com

<sup>2</sup>Department of Mathematics, Kuvempu University,  
Shankaraghatta-577451, Shimoga, Karnataka, India

<sup>3</sup>Department of Mathematics, Kuvempu University,  
Shankaraghatta-577451, Shimoga, Karnataka, India

**Abstract.** The purpose of the paper is to give some relation between the original Finslerian hypersurface and other C-conformal Finslerian hypersurfaces. In this paper we define three types of hypersurfaces, which were called a hyperplane of the 1<sup>st</sup> kind, hyperplane of the 2<sup>nd</sup> kind and hyperplane of the 3<sup>rd</sup> kind under consideration of C-conformal metric transformation.

*Key words:* Finsler spaces, Finsler hypersurface, Conformal, C-conformal, Hyperplane of 1<sup>st</sup> kind, 2<sup>nd</sup> kind and 3<sup>rd</sup> kind.

**Abstrak.** Tujuan dari paper ini adalah untuk memberikan beberapa kaitan antara hypersurface Finsler asal dengan hypersurfaces C-konformal Finsler yang lain. Dalam tulisan ini kami mendefinisikan tiga jenis hypersurfaces, yang disebut hyperplane jenis pertama, hyperplane jenis kedua dan hyperplane jenis ketiga berdasarkan transformasi metrik C-konformal.

*Kata kunci:* Ruang Finsler, hypersurface Finsler, konformal, C-konformal, hyperplane jenis pertama, jenis kedua dan jenis ketiga.

### 1. Introduction

The conformal theory and its related concepts of Finsler spaces was initiated by Knebelman in 1929. M. Hashiguchi [1] introduced a special change called C-conformal change which satisfies C-condition. The theory of Special Finsler spaces and their properties were studied by M. Matsumoto [8], C. Shibata [13] et al and authors like H. Izumi [2], S. Kikuchi [4] et al have given the condition for Finsler space to be conformally flat. C. Shibata and H. Azuma [13] have studied C-conformal

---

2000 Mathematics Subject Classification: 53C60.

Received: 05-08-2011, revised: 08-08-2011, accepted: 08-08-2011.

invariant tensor of Finsler metric. The author M. Kitayama ([5], [6], [7]) have studied Finsler spaces admitting a parallel vector field and also studied Finslerian hypersurface and metric transformations. The authors H.G. Nagaraja, C.S. Bagewadi and H. Izumi [9] have published a paper on infinitesimal h-conformal motions of Finsler metric.

The authors S.K. Narasimhamurthy and C.S. Bagewadi ([10], [11]) have published a paper on C-conformal Special Finsler spaces admitting a parallel vector field and the same authors have also studied on Infinitesimal C-conformal motions of special Finsler spaces.

Throughout the paper, terminology and notations are referred to [1], [8] and [12].

## 2. Preliminaries

A Finsler space, we mean a triple  $F^n = (M, D, L)$ , where  $M$  denotes  $n$ -dimensional differentiable manifold,  $D$  is an open subset of a tangent vector bundle  $TM$  endowed with the differentiable structure induced by the differentiable manifold  $TM$  and  $L : D \rightarrow R$  is a differentiable mapping having the properties

- i)  $L(x, y) > 0$ , for  $(x, y) \in D$ ,
- ii)  $L(x, \lambda y) = |\lambda|L(x, y)$ , for any  $(x, y) \in D$  and  $\lambda \in R$ , such that  $(x, \lambda y) \in D$ ,
- iii)  $g_{ij}(x, y) = \frac{1}{2}\dot{\partial}_i\dot{\partial}_jL^2$ ,  $(x, y) \in D$ , is positive definite, where  $\dot{\partial}_i = \frac{\partial}{\partial y^i}$ .

The metric tensor  $g_{ij}(x, y)$  and Cartan's C-tensor  $C_{ijk}$  are given by [12]:

$$g_{ij}(x, y) = \frac{1}{2}\dot{\partial}_i\dot{\partial}_jL^2, \quad g^{ij} = (g_{ij})^{-1},$$

$$C_{ijk} = \frac{1}{2}\dot{\partial}_k g_{ij}, \quad C_{jk}^i = \frac{1}{2}g^{im}(\dot{\partial}_k g_{mj}),$$

where  $\dot{\partial}_j = \frac{\partial}{\partial y^j}$  and  $\dot{\partial}_i = \frac{\partial}{\partial x^i}$ . We use the following [12]:

- a)  $l_i = \dot{\partial}_i L$ ,  $l^i = y^i/L$ ,  $h_{ij} = g_{ij} - l_i l_j$ ,
- b)  $\gamma_{jk}^i = \frac{1}{2}g^{ir}(\partial_j g_{rk} + \partial_k g_{rj} + \partial_r g_{jk})$ ,
- c)  $G^i = \frac{1}{2}\gamma_{jk}^i y^j y^k$ ,  $G_j^i = \dot{\partial}_j G^i$ ,  $G_{jk}^i = \dot{\partial}_k G_j^i$ ,  $G_{jkl}^i = \dot{\partial}_l G_{jk}^i$ , (1)
- d)  $F_{jk}^i = \frac{1}{2}g^{ir}(\delta_j g_{rk} + \delta_k g_{rj} - \delta_r g_{jk})$ ,
- e)  $N_j^i = N_j^i - y_j \sigma^i + \sigma_0 \delta_j^i + \sigma_j y^i$ ,

where  $\delta_j = \partial_j - G_j^r \partial_r$ .

The Berwald connection and the Cartan connection of  $F^n$  are given by  $B\Gamma = (G_{jk}^i, N_j^i, 0)$  and  $C\Gamma = (F_{jk}^i, N_j^i, C_{jk}^i)$  respectively.

A hypersurface  $M^{n-1}$  of the underlying smooth manifold  $M^n$  may be parametrically represented by the equation

$$x^i = x^i(u^\alpha),$$

where  $u^\alpha$  are Gaussian coordinates on  $M^{n-1}$  and Greek indices take values 1 to  $n-1$ . Here we shall assume that the matrix consisting of the projection factors  $B_\alpha^i = \partial x^i / \partial u^\alpha$  is of rank  $(n-1)$ . The following notations are also employed [6]:

$$B_{\alpha\beta}^i = \partial x^i / \partial u^\alpha \partial u^\beta, \quad B_{0\beta}^i = v^\alpha B_{\alpha\beta}^i, \quad B_{\alpha\beta\dots}^{ij\dots} = B_\alpha^i B_\beta^j \dots$$

If the supporting element  $y^i$  at a point  $(u^\alpha)$  of  $M^{n-1}$  is assumed to be tangential to  $M^{n-1}$ , we may then write

$$y^i = B_\alpha^i(u) v^\alpha,$$

i.e.,  $v^\alpha$  is thought of as the supporting element of  $M^{n-1}$  at a point  $(u^\alpha)$ . Since the function  $\underline{L}(u, v) = L(x(u), y(u, v))$  gives rise to a Finsler matrix of  $M^{n-1}$ , we get a  $(n-1)$ -dimensional Finsler space  $F^{n-1} = (M^{n-1}, \underline{L}(u, v))$ .

At each point  $(u^\alpha)$  of  $F^{n-1}$ , the unit normal vector  $N^i(u, v)$  is defined by

$$g_{ij} B_\alpha^i N^j = 0, \quad g_{ij} N^i N^j = 1. \quad (2)$$

If  $(B_\alpha^i, N_i)$  is the inverse matrix of  $(B_i^\alpha, N^i)$ , we have

$$B_\alpha^i B_i^\beta = \delta_\alpha^\beta, \quad B_\alpha^i N_i = 0, \quad N^i B_i^\alpha = 0, \quad N^i N_i = 1,$$

and further

$$B_\alpha^i B_j^\alpha + N^i N_j = \delta_j^i.$$

Making use of the inverse  $(g^{\alpha\beta})$  of  $(g_{\alpha\beta})$ , we get

$$B_i^\alpha = g^{\alpha\beta} g_{ij} B_\beta^j, \quad N_i = g_{ij} N^j.$$

For the induced Cartan connections  $ICT = (F_{\beta\gamma}^\alpha, N_\beta^\alpha, C_{\beta\gamma}^\alpha)$  on  $F^{n-1}$ , the second fundamental h-tensor  $H_{\alpha\beta}$  and the normal curvature tensor  $H_\alpha$  are given by

$$i) \quad H_{\alpha\beta} = N_i (B_{\alpha\beta}^i + F_{jk}^i B_{\alpha\beta}^{jk}) + M_\alpha H_\beta, \quad (3)$$

$$ii) \quad H_\alpha = N_i (B_{0\alpha}^i + N_j^i B_\alpha^j),$$

respectively, where  $M_\alpha = C_{ijk} B_\alpha^i N^j N^k$  and  $B_{0\alpha}^i = B_{\beta\alpha}^i v^\beta$ . Transvecting  $H_{\alpha\beta}$  by  $v^\beta$ , we get  $H_{0\alpha} = H_{\beta\alpha} v^\beta = H_\alpha$ .

Further more we have to put

$$M_{\alpha\beta} = C_{ijk} B_{\alpha\beta}^{ij} N^k. \quad (4)$$

### 3. C-Conformal Finsler Space

We shall consider conformal change of a Finsler metric formed by  $L \rightarrow \bar{L} = e^{\sigma(x)}L$ , where  $\sigma$  is conformal factor depends on the point  $x$  only and under this change we have another Finsler space  $\bar{F}^n = (M^n, \bar{L})$  on the same underlying manifold  $M^n$ .

M. Hashiguchi [1] introduced the special change named C-conformal change which is by definition, a non-homothetic conformal change satisfying

$$C_{jk}^i \sigma^j = 0, \quad (5)$$

where  $C_{jk}^i = g^{im}(\partial_j g_{km})/2$ ,  $\sigma^i = g^{im}\sigma_m$ ,  $\sigma_m = \partial\sigma/\partial x^m$ ,  $\sigma^j = g^{ij}\sigma_j$ . From (1) and by symmetry of lower indices of  $C_{ijk}$ , we have

$$C_{ijk}\sigma^i = C_{jik}\sigma^i = C_{jki}\sigma^i = 0,$$

also we have

$$C_{ij}^k \sigma^i = C_{ij}^k \sigma^j = C_{jk}^i \sigma^k = 0.$$

In the following the quantity with bar will be defined in C-conformal Finsler space  $\bar{F}^n$ , and the quantity without bar will be defined in Finsler space  $F^n$ . Under the C-conformal change, we have the following [2], [13]:

$$\begin{aligned} a) \quad & \bar{g}_{ij} = (\bar{L}/L)^2 g_{ij}, \quad \bar{g}^{ij} = (L/\bar{L})^2 g^{ij}, \\ b) \quad & \bar{y}_i = (\bar{L}/L)^2 y_i, \\ c) \quad & \bar{C}_{ijk} = C_{ijk}, \quad \bar{C}_{jk}^i = e^{2\sigma} C_{jk}^i, \quad \bar{C}_i = e^{-2\sigma} C_i, \\ d) \quad & \bar{\gamma}_{jk}^i = \gamma_{jk}^i + (\sigma_j \delta_k^i + \sigma_k \delta_j^i - g_{jk} \sigma^i), \\ e) \quad & \bar{G}^i = G^i - \frac{1}{2} L^2 \sigma^i + \sigma_0 y^i, \\ f) \quad & \bar{G}_{jk}^i = G_{jk}^i - g_{jk} \sigma^i + \sigma_k \delta_j^i + \sigma_j \delta_k^i, \\ g) \quad & \bar{N}_j^i = N_j^i - y_j \sigma^i + \sigma_0 \delta_j^i + \sigma_j y^i, \\ h) \quad & \bar{F}_{jk}^i = F_{jk}^i - g_{jk} \sigma^i + \sigma_k \delta_j^i + \sigma_j \delta_k^i + \sigma_0 C_{jk}^i. \end{aligned} \quad (6)$$

### 4. Hypersurface Given by a C-Conformal Change

We now consider a Finsler hypersurface  $F^{n-1} = (M^{n-1}, \underline{L}(u, v))$  of the Finsler space  $F^n$  and another Finsler hypersurface  $\bar{F}^{n-1} = (M^{n-1}, \underline{\bar{L}}(u, v))$  of the Finsler space  $F^n$  given by the C-conformal change.

Let  $N^i(u, v)$  be a unit normal vector at each point of the  $F^{n-1}$ , and as component of  $n-1$  linearly independent tangent vectors of  $F^{n-1}$  and they are invariant under the C-conformal change. Thus we shall show that a unit normal vector  $\bar{N}^i(u, v)$  of  $\bar{F}^{n-1}$  is uniquely determined by

$$\bar{g}_{ij} B_\alpha^i \bar{N}^j = 0, \quad \bar{g}_{ij} \bar{N}^i \bar{N}^j = 1. \quad (7)$$

By means of (2) and (6), we get

$$\bar{g}_{ij}(\pm e^{-\sigma} N^i)(\pm e^{-\sigma} N^j) = 1.$$

Therefore we can put

$$\bar{N}^i = e^{-\sigma} N^i,$$

where we have chosen the sign '+' in order to fix an orientation. It is obvious that  $\bar{N}_i(u, v)$  satisfies (2), hence we obtain:

**Lemma 4.1.** *For a field of linear frame  $(B_1^i, B_2^i, \dots, B_{n-1}^i, N^i)$  of  $F^n$ , there exists a field of linear frame  $(B_1^i, B_2^i, \dots, B_{n-1}^i, \bar{N}^i = e^{-\sigma} N^i)$  of the  $\bar{F}^n$  given by the C-conformal change such that (7) satisfied along  $\bar{F}^{n-1}$ .*

The quantities  $\bar{B}_i^\alpha$  are uniquely defined along  $\bar{F}^{n-1}$  by

$$\bar{B}_i^\alpha = \bar{g}^{\alpha\beta} \bar{g}_{ij} B_j^\beta,$$

where  $(\bar{g}^{\alpha\beta})$  is the inverse metric of  $(\bar{g}_{\alpha\beta})$ . If  $(\bar{B}_i^\alpha, \bar{N}^i)$  is the inverse vector of  $(\bar{B}_\alpha^i, \bar{N}_i)$ , then we have

$$B_\alpha^i \bar{B}_i^\beta = \delta_\alpha^\beta, \quad B_\alpha^i \bar{N}_i = 0, \quad \bar{N}^i \bar{B}_i^\alpha = 0, \quad \bar{N}^i \bar{N}_i = 1,$$

and also

$$B_\alpha^i \bar{B}_j^\alpha + \bar{N}^i \bar{N}_j = \delta_j^i.$$

Also we get  $\bar{N}_i = \bar{g}_{ij} \bar{N}^j$ , that is

$$\bar{N}_i = e^\sigma N_i. \quad (8)$$

We have from (6(e)),

$$D^i = \bar{G}^i - G^i = \sigma_0 y^i - \frac{L^2}{2} \sigma^i, \quad \text{where } \sigma_0 = \sigma_r y^r. \quad (9)$$

Differentiating (9) by  $y^j$  and from (6(f)), we obtain

$$\begin{aligned} D_j^i &= D_{(j)}^i, \\ &= \bar{G}_j^i - G_j^i, \\ &= \bar{N}_j^i - N_j^i, \\ &= -y_j \sigma^i + \sigma_0 \delta_j^i + \sigma_j y^i, \end{aligned}$$

where  $D_{(j)}^i = \dot{\partial}_j D^i$ . From (9), we have

$$N_i D^i = \sigma_0 N_i y^i - \frac{L^2}{2} N_i \sigma^i.$$

We assume that  $N_i \sigma^i = 0$ . i.e.,  $\sigma^i(x)$  is tangential to  $F^{n-1}$  and using the condition  $N_i y^i = 0$ , then we have

$$N_i D^i = 0. \quad (10)$$

Differentiating (10) by  $y^j$ , we have

$$\begin{aligned} N_i D_{(j)}^i + D^i (N_i)_{(j)} &= 0, \\ N_i D_j^i + D^i (\partial_j N_i) &= 0. \end{aligned}$$

Transvecting above equation by  $B_\alpha^j$ , we get

$$\begin{aligned} N_i D_j^i B_\alpha^j + D^i (\partial_j N_i) B_\alpha^j &= 0, \\ N_i D_j^i B_\alpha^j &= 0, \end{aligned} \tag{11}$$

where we used

$$B_\alpha^j (\partial_j N_i) = M_\alpha N_i = C_{ijk} B_\alpha^j N^i N^k N_i = 0.$$

**Definition 4.1.** *If each path of the hypersurface  $F^{n-1}$  with respect to the induced connection is also a path of the ambient space  $F^n$ , then  $F^{n-1}$  is called a 'hyperplane of the 1<sup>st</sup> kind'.*

A hyperplane of the 1<sup>st</sup> kind is characterized by  $H_\alpha = 0$ .

From (3(ii)) and using (8), we have

$$\bar{H}_\alpha = \bar{N}_i (B_{0\alpha}^i + \bar{N}_j^i B_\alpha^j).$$

Thus

$$\begin{aligned} \bar{H}_\alpha - e^\sigma H_\alpha &= \bar{N}_i (B_{0\alpha}^i + \bar{N}_j^i B_\alpha^j) - e^\sigma N_i (B_{0\alpha}^i + N_j^i B_\alpha^j), \\ &= e^\sigma (N_i B_{0\alpha}^i + N_i \bar{N}_j^i B_\alpha^j) - e^\sigma (N_i B_{0\alpha}^i + N_i N_j^i B_\alpha^j), \\ &= e^\sigma N_i (\bar{N}_j^i - N_j^i) B_\alpha^j, \\ &= e^\sigma N_i D_j^i B_\alpha^j. \end{aligned}$$

Thus we have

$$\bar{H}_\alpha = e^\sigma (H_\alpha + N_i D_j^i B_\alpha^j).$$

Thus from (11), we obtained

$$\bar{H}_\alpha = e^\sigma H_\alpha.$$

Hence we state the following:

**Theorem 4.1.** *A Finsler hypersurface  $F^{n-1}$  is a hyperplane of 1<sup>st</sup> kind if and only if  $C$ -conformal Finsler hypersurface  $\bar{F}^{n-1}$  is a hyperplane of 1<sup>st</sup> kind, provided  $N_i \sigma^i = 0$ , i.e.,  $\sigma^i(x)$  is tangential to  $F^{n-1}$ .*

Now from (6(h)), the so called difference tensor  $D_{jk}^i$  has the following form

$$\begin{aligned} D_{jk}^i &= \bar{F}_{jk}^i - F_{jk}^i, \\ &= -g_{ij} \sigma^i + \sigma_k \delta_j^i + \sigma_j \delta_k^i + \sigma_0 C_{jk}^i. \end{aligned}$$

Contracting above equation by  $N_i$  and  $B_\alpha^j$ , we get

$$\begin{aligned} N_i D_{jk}^i B_\alpha^j &= -N_i g_{jk} \sigma^i B_\alpha^j + \sigma_k N_i \delta_j^i B_\alpha^j + \sigma_j N_i \delta_k^i B_\alpha^j + \sigma_0 C_{jk}^i N_i B_\alpha^j, \\ &= 0. \end{aligned}$$

Where we use  $\sigma_0 = \sigma_i y^i$  and equation (5). Thus we state the following:

**Lemma 4.2.** *Assuming that  $\sigma_i(x)$  is tangential to  $F^{n-1}$ , then the tensor  $N_i D_{jk}^i B_{\alpha}^j$  is vanishes if and only if it satisfies (5).*

**Definition 4.2.** *If each  $h$ -path of a hypersurface  $F^{n-1}$  with respect to the induced connection is also  $h$ -path of the ambient space  $F^n$ , then  $F^{n-1}$  is called a ‘hyperplane of the 2<sup>nd</sup> kind’.*

A hyperplane of the 2<sup>nd</sup> kind is characterized by  $H_{\alpha\beta} = 0$ .

From (3(i)), we have

$$H_{\alpha\beta} = N_i(B_{\alpha\beta}^i + F_{jk}^i B_{\alpha\beta}^{jk}) + M_{\alpha} H_{\beta}. \quad (12)$$

Under the C-conformal change, (12) can be written as

$$\bar{H}_{\alpha\beta} = \bar{N}_i(B_{\alpha\beta}^i + \bar{F}_{jk}^i B_{\alpha\beta}^{jk}) + \bar{M}_{\alpha} \bar{H}_{\beta}. \quad (13)$$

Using equations (12) and (13), we get

$$\begin{aligned} \bar{H}_{\alpha\beta} - e^{\sigma} H_{\alpha\beta} &= [\bar{N}_i(B_{\alpha\beta}^i + \bar{F}_{jk}^i B_{\alpha\beta}^{jk}) + \bar{M}_{\alpha} \bar{H}_{\beta}] \\ &\quad - e^{\sigma} [N_i(B_{\alpha\beta}^i + F_{jk}^i B_{\alpha\beta}^{jk}) + M_{\alpha} H_{\beta}], \end{aligned} \quad (14)$$

using  $\bar{M}_{\alpha} = M_{\alpha}$  and  $\bar{H}_{\alpha} = e^{\sigma} H_{\alpha}$ , we have

$$\bar{H}_{\alpha\beta} - e^{\sigma} H_{\alpha\beta} = e^{\sigma} N_i (\bar{F}_{jk}^i - F_{jk}^i) B_{\alpha\beta}^{jk},$$

that implies

$$\bar{H}_{\alpha\beta} - e^{\sigma} H_{\alpha\beta} = e^{\sigma} (N_i D_{jk}^i B_{\alpha\beta}^{jk}). \quad (15)$$

Thus by virtue of lemma (4.1), therefore we state the following:

**Theorem 4.2.** *A Finsler hypersurface  $F^{n-1}$  is a hyperplane of the 2<sup>nd</sup> kind if and only if the C-conformal Finsler hypersurface  $\bar{F}^{n-1}$  is a hyperplane of the 2<sup>nd</sup> kind, provided  $\sigma_i(x)$  is tangential to  $F^{n-1}$ .*

**Definition 4.3.** *If the unit normal vector of  $F^{n-1}$  is parallel along each curve of  $F^{n-1}$ , then  $F^{n-1}$  is called a ‘hyperplane of the 3<sup>rd</sup> kind’.*

A hyperplane of the 3<sup>rd</sup> kind is characterized by  $H_{\alpha\beta} = M_{\alpha\beta} = 0$ .

From (4), under C-conformal change the tensor  $M_{\alpha\beta}$  can be written as

$$\begin{aligned} \bar{M}_{\alpha\beta} &= \bar{C}_{ijk} \bar{B}_{\alpha\beta}^{ij} \bar{N}^k, \\ &= e^{-\sigma} C_{ijk} B_{\alpha\beta}^{ij} N^k, \\ &= e^{-\sigma} M_{\alpha\beta}. \end{aligned} \quad (16)$$

By characterization of hyperplane of the 3<sup>rd</sup> kind and (15), we have  $\bar{H}_{\alpha\beta} = \bar{M}_{\alpha\beta} = 0$ .

Thus by virtue of lemma (4.1), we state the following:

**Theorem 4.3.** *A Finsler hypersurface  $F^{n-1}$  is a hyperplane of the 3<sup>rd</sup> kind if and only if C-conformal Finsler hypersurface  $\bar{F}^{n-1}$  is a hyperplane of the 3<sup>rd</sup> kind, provided  $\sigma_i(x)$  is tangential to  $F^{n-1}$ .*

**Acknowledgement.** The authors are thankful to the referees for their valuable suggestions.

### References

- [1] Hashiguchi, M., "On conformal transformations of Finsler metric", *J. Math. Kyoto Univ.*, **16** (1976), 25-50.
- [2] Izumi, H., "Conformal transformations of Finsler spaces", *Tensor, N.S.*, **34** (1980), 337-359.
- [3] Zilasi, J., and Vincze, C., "A new look at Finsler connections and special Finsler manifolds", *Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis*, **16** (2000), 33-63.
- [4] Kikuchi, S., "On condition that a Finsler space be conformally flat", *Tensor, N.S.*, **55** (1994), 97-100.
- [5] Kitayama, M., "Finsler spaces admitting a parallel vector field", *Balkan J. of Geometry and its Applications*, **3** (1998), 29-36.
- [6] Kitayama, M., "Finslerian hypersurfaces and metric transformations", *Tensor, N.S.*, **60** (1998), 171-177.
- [7] Kitayama, M., "On Finslerian hypersurfaces given by  $\beta$  change", *Balkan J. of Geometry and its Applications*, **7** (2002), 49-55.
- [8] Matsumoto, M., *Foundations of Finsler geometry and special Finsler spaces*, Kaiseisha press, Otsu, Saikawa, 1986.
- [9] Nagaraja, H.G., and Bagewadi, C.S., "Finsler spaces admitting semi-symmetric Finsler connections", *Advances in modelling and Analysis*, **A** (2003), 57-68.
- [10] Narasimhamurthy, S.K., Bagewadi, C.S., and Nagaraja, H.G., "Infinitesimal C-conformal motions of special Finsler spaces", *Tensor, N.S.*, **64** (2003), 241-247.
- [11] Narasimhamurthy, S.K., and Bagewadi, C.S., "C-conformal Finsler spaces admitting a parallel vector field", *Tensor, N.S.*, **65** (2004), 162-169.
- [12] Rund, H., *The differential geometry of Finsler spaces*, Springer-Verlag, Berlin, 1959.
- [13] Shibata, C., and Azuma, M., "C-conformal invariant tensors of Finsler metrics", *Tensor, N.S.*, **52** (1993), 76-81.